arXiv:1207.3853v1 [math.DG] 17 Jul 2012

INTRINSIC INVARIANTS OF CROSS CAPS

M. HASEGAWA, A. HONDA, K. NAOKAWA, M. UMEHARA, AND K. YAMADA

Abstract. It is classically known that generic smooth maps of $\boldsymbol{R}^2$ into $\boldsymbol{R}^3$ admit only cross cap singularities. This suggests that the class of cross caps might be an important object in differential geometry. We show that the *standard cross cap* $f_{\mathrm{std}}(u,v)=(u,uv,v^2)$ has non-trivial isometric deformations with infinite dimensional freedom. Since there are several geometric invariants for cross caps, the existence of isometric deformations suggests that one can ask which invariants of cross caps are intrinsic. In this paper, we show that there are three fundamental intrinsic invariants for cross caps. The existence of extrinsic invariants is also shown.

Introduction

Let U be a domain in $\boldsymbol{R}^2$ and $f : U \to \boldsymbol{R}^3$ a C^∞-map. A point $p\ (\in U)$ is called a *singular point* if the rank of the Jacobi matrix of f at p is less than 2. Consider such a map given by

$$f_{\mathrm{std}}(u,v)=(u,uv,v^2), \tag{1}$$

which has an isolated singular point at the origin $(0,0)$ and is called the *standard cross cap* (see Figure 1, left). A singular point p of a map $f : U \to \boldsymbol{R}^3$ is called a *cross cap* or a *Whitney umbrella* if there exist local diffeomorphism φ on $\boldsymbol{R}^2$ and a local diffeomorphism Φ on $\boldsymbol{R}^3$ such that $\Phi\circ f = f_{\mathrm{std}}\circ\varphi$. Whitney proved that a C^∞-map $f : U \to \boldsymbol{R}^3$ has a cross cap singularity at $p \in U$ if there exists a local coordinate system (u,v) centered at p such that

$$f_v(0,0) := \frac{\partial f}{\partial v}(0,0) = 0$$

and three vectors

$$f_u(0,0) := \frac{\partial f}{\partial u}(0,0), \quad f_{uv}(0,0) := \frac{\partial^2 f}{\partial u \partial v}(0,0), \quad f_{vv}(0,0) := \frac{\partial^2 f}{\partial v^2}(0,0)$$

Date: July 17, 2012.

2010 *Mathematics Subject Classification.* Primary 57R45; Secondary 53A05.

Key words and phrases. cross cap, curvature, isometric deformation .

The second and third authors were partly supported by the Grant-in-Aid for JSPS Fellows. The fourth and fifth authors were partially supported by Grant-in-Aid for Scientific Research (A) No. 22244006, and Scientific Research (B) No. 21340016, respectively, from the Japan Society for the Promotion of Science.



FIGURE 1. An isometric deformation of the standard cross cap

are linearly independent. By a rotation, a translation in $\boldsymbol{R}^3$ and a suitable orientation preserving coordinate change of the domain $U \subset \boldsymbol{R}^2$, we have the following Maclaurin expansion of f at a cross cap singularity $(0,0)$ (cf. [12] or [3])

$$f(u,v) = \left(u, uv + \sum_{i=3}^{n} \frac{b_i}{i!} v^i, \sum_{r=2}^{n} \sum_{j=0}^{r} \frac{a_{j\,r-j}}{j!(r-j)!} u^j v^{r-j} \right) + O(u,v)^{n+1}, \tag{2}$$

where a_{02} never vanishes. By orientation preserving coordinate changes $(u,v) \mapsto (-u,-v)$ and $(x,y,z) \mapsto (-x,y,-z)$, we may assume that

$$a_{02} > 0, \tag{3}$$

where (x,y,z) is the usual Cartesian coordinate system of $\boldsymbol{R}^3$. After this normalization (3), one can easily verify that all of the coefficients a_{jk} and b_i are uniquely determined. An oriented local coordinate system (u,v) giving such a normal form is called the *canonical coordinate system* of f at the cross cap singularity. This unique expansion of a cross cap implies that the coefficients a_{jk} and b_i can be considered as geometric invariants of the cross cap f. A cross cap is called *non-degenerate* (resp. *degenerate*) if a_{20} does not vanish (resp. does vanish). On the other hand, a cross cap is called *quadratic* if $a_{jk} = 0$ for $j+k \geq 3$ and $b_i = 0$ for $i \geq 3$. The standard cross cap is a typical example of a degenerate quadratic cross cap.

Let f be a degenerate quadratic cross cap. The first main result of this paper is to show the existence of non-trivial isometric deformations of f (see Figure 1). More precisely, we show that each spherical open regular arc with a base point induces a cross cap, not congruent to f, but with first fundamental form equal to that of f. Since the congruent classes appearing in such an isometric deformation can be distinguished by the geodesic curvature functions of the corresponding spherical curves, this implies that each degenerate quadratic cross cap induces a non-trivial family of isometric deformations with infinite dimensional freedom. Moreover, using such a deformation, we show that the invariants a_{03}, a_{12} and b_3 in (2) are extrinsic, namely, these invariants change according to the isometric deformation.

The differential geometry of cross caps in $\boldsymbol{R}^3$ has been discussed by several authors (cf. [1], [2], [3], [4], [5], [6], [8], [11] and [12]). However, the distinction between intrinsic and extrinsic invariants has not been clearly discussed before. When $f : U \to \boldsymbol{R}^3$ is an immersion, then it induces a Riemannian metric on U (called the first fundamental form) and we know that 'intrinsic' means that a given invariant is described in terms of this Riemannian structure on U. Similarly, each cross cap induces a positive semidefinite symmetric tensor ds^2 as a pull-back of the ambient metric. Then a given invariant of a cross cap is called *intrinsic* if it can be described in terms of this positive semidefinite metric ds^2. In the case of cuspidal edges in $\boldsymbol{R}^3$, such an intrinsic invariant is defined as a 'singular curvature' along these singular points (cf. [9]).

We then show that a_{02}, a_{20} and a_{11} are intrinsic invariants. In fact, if a_{20} is negative then the Gaussian curvature of a given cross cap is negative and having no lower bound. On the other hand, if a_{20} is positive, then the Gaussian curvature is not bounded from below nor from above. Fukui and the first author [3] found an important concept of the 'focal conic' of a cross cap, as a section of its caustic by the normal plane (see the explanation after (15) and also [4]). They also showed that focal conics have the expression

$$y^2 + 2a_{11}yz - (a_{20}a_{02} - a_{11}^2)z^2 + a_{02}z = 0.$$

The focal conic is a hyperbola (resp. an ellipse) if and only if a_{20} is positive (resp. negative). Since we have seen that a_{02}, a_{20} and a_{11} are intrinsic, we can say that focal conics live in the intrinsic geometry of cross caps, although caustics themselves are extrinsic objects. It should be remarked that the Gauss-Bonnet type formula for closed surfaces which admit only cross cap singularities in $\boldsymbol{R}^3$ has no defection at each singular point (cf. Kuiper [7, p. 92]).

Isometric deformations of degenerate quadratic cross caps

As mentioned in the introduction, a quadratic cross cap can be expressed as

$$\left(u, uv, \frac{1}{2}(a_{20}u^2 + 2a_{11}uv + a_{02}v^2)\right) \qquad (a_{02} > 0),$$

where (u, v) is the canonical coordinate system. As defined in the introduction, the cross cap $(0, 0)$ is degenerate if a_{20} vanishes. A degenerate quadratic cross cap has the following expression

$$f_0(u, v) := \frac{1}{2}\left(0, 0, a_{02}v^2\right) + u\,(1, v, a_{11}v) \qquad (a_{02} > 0). \tag{4}$$

In particular, it is a ruled surface. The first fundamental form

$$ds^2 := E_0\,du^2 + 2F_0\,du\,dv + G_0\,dv^2$$

of f_0 is given by

$$
\begin{aligned}
E_0 &:= (f_0)_u \cdot (f_0)_u = 1 + (1 + a_{11}^2)v^2, \\
F_0 &:= (f_0)_u \cdot (f_0)_v = (1 + a_{11}^2)uv + a_{02}a_{11}v^2, \\
G_0 &:= (f_0)_v \cdot (f_0)_v = (1 + a_{11}^2)u^2 + 2a_{02}a_{11}uv + a_{02}^2 v^2,
\end{aligned}
$$

where the dot indicates the canonical inner product of $\boldsymbol{R}^3$.

Definition 1. Let U be a domain in $(\boldsymbol{R}^2; u, v)$ containing the origin, and let $f_i : U \to \boldsymbol{R}^3$ $(i = 0, 1)$ be two C^∞-maps having a cross cap singularity at $(0, 0)$. If f_0 and f_1 satisfy

$$
(f_0)_u \cdot (f_0)_u = (f_1)_u \cdot (f_1)_u, \quad (f_0)_u \cdot (f_0)_v = (f_1)_u \cdot (f_1)_v, \\
(f_0)_v \cdot (f_0)_v = (f_1)_v \cdot (f_1)_v,
$$

then we say that f_0 is *isometric* to f_1. On the other hand, let $f_t : U \to \boldsymbol{R}^3$ $(|t| < \epsilon)$, be a smooth 1-parameter family of C^∞-maps having a cross cap singularity at $(0, 0)$, where ϵ is a positive constant. Then $\{f_t\}_{|t|<\epsilon}$ is called an *isometric deformation* if each f_t is isometric to f_0. An isometric deformation of f_t is *non-trivial* if each f_t is not congruent to f_0.

We prove the following assertion.

Theorem 2. *Let $c(s)$ $(|s| < \pi/2)$ be a regular curve in the unit sphere $S^2 (\subset \boldsymbol{R}^3)$ with arc length parameter. We set*

$$
\xi(v) := \sqrt{1 + (1 + a_{11}^2)v^2}\, \hat{c}(v), \qquad \hat{c}(v) := c\left(\arctan\left(v\sqrt{1 + a_{11}^2}\right)\right), \tag{5}
$$

for each $v \in \boldsymbol{R}$, and

$$
\gamma(v) := \frac{a_{02}}{1 + a_{11}^2} \int_0^v tB(t)dt, \qquad B(v) := a_{11}\xi'(v) + \xi(v) \times \xi'(v),
$$

where the prime means the derivative with respect to v and $\times$ denotes the vector product in $\boldsymbol{R}^3$. Then a ruled surface $f_c : \boldsymbol{R}^2 \to \boldsymbol{R}^3$ defined by

$$
f_c(u, v) := \gamma(v) + u\xi(v)
$$

has a cross cap singularity at the origin such that f_c is isometric to a degenerate quadratic cross cap f_0. Moreover, let $c_i(s)$ $(|s| < \pi/2;\ i = 1, 2)$ be two regular curves in S^2 with arc length parameter. Then f_{c_1} is congruent to f_{c_2} if and only if c_1 is congruent to c_2 in S^2. In particular, f_c is congruent to f_0 as in (4) if and only if $c(s)$ is a geodesic in S^2.

Proof. By (5), we have that

$$
\xi(v) \cdot \xi(v) = 1 + (1 + a_{11}^2)v^2, \tag{6}
$$

$$
\xi'(v) \cdot \xi'(v) = 1 + a_{11}^2, \tag{7}
$$

$$
\xi(v) \cdot \xi'(v) = (1 + a_{11}^2)v, \tag{8}
$$

where we used the fact that $\hat{c}'(v)$ is orthogonal to $\hat{c}(v)$. Since $\xi(v)\times\xi'(v)$ is orthogonal to $\xi(v)$ and $\xi'(v)$, the equations (7) and (8) yield that

$$\xi(v)\cdot B(v) = a_{11}(1+a_{11}^2)v, \qquad \xi'(v)\cdot B(v) = a_{11}(1+a_{11}^2).$$

Finally, we have

$$B\cdot B = |\xi\times\xi'|^2 + a_{11}^2|\xi'|^2 = |\xi|^2|\xi'|^2 - (\xi\cdot\xi')^2 + a_{11}^2|\xi'|^2 = (1+a_{11}^2)^2.$$

From now on, we denote $f := f_c$ for the sake of simplicity. One can prove that $f_u\cdot f_u$, $f_u\cdot f_v$ and $f_v\cdot f_v$ coincide with E_0, F_0 and G_0 respectively, using the above relations.

The unit normal vector field $\nu(u,v)$ is given by

$$\nu(u,v) = \frac{1}{\delta}\biggl(-\Bigl(a_{02}v\sqrt{1+(1+a_{11}^2)v^2}\Bigr)\boldsymbol{e}(v) + \bigl((1+a_{11}^2)u + a_{02}a_{11}v\bigr)\boldsymbol{n}(v)\biggr),$$

where

$$\delta := \sqrt{1+a_{11}^2}\sqrt{(1+a_{11}^2)u^2 + 2a_{02}a_{11}uv + a_{02}^2v^2(1+v^2)},$$

$\boldsymbol{e} = dc/ds$ and $\boldsymbol{n} = c\times\boldsymbol{e}$. By a straightforward calculation, the second fundamental form $L\,du^2 + 2M\,du\,dv + N\,dv^2$ of f is given by

$$\begin{aligned} &L := 0, \quad M := -\frac{a_{02}\sqrt{1+a_{11}^2}v}{\delta}, \\ &N := \frac{a_{02}\sqrt{1+a_{11}^2}}{\delta}u + \frac{\delta\kappa(v)}{(1+(1+a_{11}^2)v^2)^{3/2}}, \end{aligned} \tag{9}$$

where $\kappa(s)$ is the geodesic curvature of $c(s)$. Let $c_i(s)$ ($|s|<\pi/2$; $i=1,2$) be two regular curves in S^2 with arc length parameter, and $\kappa_i(s)$ the geodesic curvature function of $c_i(s)$. Then f_{c_1} is congruent to f_{c_2} if c_1 is congruent to c_2 in S^2, since (9) implies that the second fundamental form of f_{c_1} coincides with that of f_{c_2} if and only if κ_1 coincides with κ_2. Finally, as seen in the following corollary, degenerate quadratic cross caps correspond to the great circles, so we get the assertion. □

Corollary 3. *There exists an infinite dimensional family of non-trivial isometric deformations of a given degenerate quadratic cross cap f_0.*

Proof. Consider a line passing through $(1,0,0)$ whose velocity vector is $(0,1,a_{11})$. Then the central projection of the line into the unit sphere gives a subarc of a great circle

$$\hat{c}_0(v) = \frac{1}{\sqrt{1+(1+a_{11}^2)v^2}}(1,v,a_{11}v).$$

The degenerate quadratic cross cap f_0 can be constructed via $\hat{c}_0$ by applying Theorem 2. Then any non-trivial deformation of the curve $\hat{c}_0$ as a spherical curve gives a non-trivial isometric deformation of f_0, which proves the assertion. □

For example, take a constant κ and set

$$c_\kappa(s) := \frac{1}{\mu^2}\Bigl(\kappa^2+\cos(\mu s), \mu\sin(\mu s), \kappa\bigl(1-\cos(\mu s)\bigr)\Bigr) \quad \left(|s|<\frac{\pi}{2},\ \mu := \sqrt{1+\kappa^2}\right),$$

which gives a circle in S^2 with arc length parameter and of constant geodesic curvature κ. Then it produces a deformation of the standard cross cap, where c_0 corresponds to f_{std} as in (1). Figure 1 indicates the cross caps corresponding to $\kappa = 0$, 1 and 3, respectively.

Using the existence of this non-trivial isometric deformation of degenerate quadratic cross caps, we can prove the following assertion.

Theorem 4. *The three invariants a_{12}, a_{03} and b_3 are extrinsic.*

Proof. Let $c(s)$ be a spherical curve with arc length parameter s so that

$$c(0) = (1,0,0), \qquad \frac{dc}{ds}(0) = \frac{1}{\sqrt{1+a_{11}^2}}(0,1,a_{11}),$$

and let $\boldsymbol{e} := dc/ds$, $\boldsymbol{n} := c \times \boldsymbol{e}$. Then

$$\frac{d\boldsymbol{e}}{ds}(s) = \kappa(s)\boldsymbol{n}(s) - c(s), \qquad \frac{d\boldsymbol{n}}{ds}(s) = -\kappa(s)\boldsymbol{e}(s),$$

hold, where $\kappa(s)$ is the curvature function. Using these, one can see that the cross cap $f = f_c$ given in Theorem 2 has the following expansion:

$$\begin{aligned}(10)\quad f(u,v) = &\left(u, uv, a_{11}uv + \frac{1}{2}a_{02}v^2\right) \\ &+ \frac{\kappa(0)\sqrt{1+a_{11}^2}}{6}\left(0, -3a_{11}uv^2 - 2a_{02}v^3, 3uv^2\right) + O(u,v)^4.\end{aligned}$$

By a parameter change $v = w + \frac{1}{2}a_{11}\kappa(0)\sqrt{1+a_{11}^2}\,w^2$, (10) is rewritten as

$$\begin{aligned}f(u,w) = &\left(u, uw, a_{11}uw + \frac{1}{2}a_{02}w^2\right) \\ &+ \frac{\kappa(0)\sqrt{1+a_{11}^2}}{6}\left(0, -2a_{02}w^3, 3(1+a_{11}^2)uw^2 + 3a_{11}a_{02}w^3\right) + O(u,w)^4.\end{aligned}$$

Thus, (u,w) forms the canonical coordinate system up to the third order terms, and the invariants a_{12}, a_{03} and b_3 are expressed as

$$a_{12} = \kappa(0)(1+a_{11}^2)^{3/2}, \quad a_{03} = 3a_{02}a_{11}\kappa(0)\sqrt{1+a_{11}^2},$$
$$b_3 = -2a_{02}\kappa(0)\sqrt{1+a_{11}^2}$$

which depend on the initial value $\kappa(0)$ of the geodesic curvature function, and thus they are extrinsic. □

Definition 5. Let $f : U \to \boldsymbol{R}^3$ be a C^∞-map and $p \in U$ a cross cap singularity. A local coordinate system (u,v) centered at p is said to be *admissible* if it satisfies $f_v(0,0) = 0$.

Canonical coordinate systems of cross caps are admissible. The concept of admissible coordinate systems is intrinsic, since $\partial/\partial v$ at $(0,0)$ points the degenerate directions of the induced metrics. In contrast to Theorem 4, the following assertion holds:

Theorem 6. *The coefficients a_{02}, a_{20} and a_{11} are intrinsic invariants of cross caps.*

Proof. Let $(0,0)$ be a cross cap singularity of a C^∞-map $f : (U; u, v) \to \boldsymbol{R}^3$, and (u, v) be an admissible coordinate system. Without loss of generality, we may assume that

$$[f_u, f_{uv}, f_{vv}] > 0$$

by applying the coordinate change $(u, v) \mapsto (-u, -v)$ if necessary, where

$$[\boldsymbol{a}, \boldsymbol{b}, \boldsymbol{c}] := \det(\boldsymbol{a}, \boldsymbol{b}, \boldsymbol{c}) = (\boldsymbol{a} \times \boldsymbol{b}) \cdot \boldsymbol{c}$$

and × is the vector product of $\boldsymbol{R}^3$.

Then we have that

$$a_{02} = \frac{|f_u|\,|f_u \times f_{vv}|^3}{[f_u, f_{uv}, f_{vv}]^2}, \tag{11}$$

$$a_{20} = \frac{|f_u \times f_{vv}|}{4|f_u|^3[f_u, f_{uv}, f_{vv}]^2}\Big([f_u, f_{uu}, f_{vv}]^2 + 4[f_u, f_{uv}, f_{vv}][f_u, f_{uv}, f_{uu}]\Big), \tag{12}$$

$$a_{11} = \frac{1}{2|f_u|[f_u, f_{uv}, f_{vv}]^2}\left(2[f_u, f_{uv}, f_{vv}] \det\begin{pmatrix} f_u \cdot f_u & f_u \cdot f_{uv} \\ f_{vv} \cdot f_u & f_{vv} \cdot f_{uv} \end{pmatrix} - |f_u \times f_{vv}|^2[f_u, f_{uu}, f_{vv}]\right) \tag{13}$$

hold at $(u, v) = (0, 0)$. One can prove these identities immediately: In fact, the right hand sides of these identities are independent of the choice of admissible coordinate systems, and these identities themselves can be directly verified for the canonical coordinate system of f.

We now set

$$E := f_u \cdot f_u, \quad F := f_u \cdot f_v, \quad G := f_v \cdot f_v,$$

which are the coefficients of the induced metric of the cross cap. It is sufficient to show that the right hand sides of (11), (12) and (13) are written in terms of derivatives of E, F and G at $(0, 0)$.

We first show that a_{02} is intrinsic: Since $f_v(0, 0) = 0$, it holds that

$$[f_u, f_{uv}, f_{vv}]^2 = \det\left(\begin{pmatrix} f_u \\ f_{uv} \\ f_{vv} \end{pmatrix} (f_u, f_{uv}, f_{vv})\right) = \det\begin{pmatrix} E & F_u & F_v \\ F_u & G_{uu}/2 & G_{uv}/2 \\ F_v & G_{uv}/2 & G_{vv}/2 \end{pmatrix} \tag{14}$$

at $(u, v) = (0, 0)$, where we used the identities

$$f_u \cdot f_u = E, \quad f_u \cdot f_{uv} = F_u, \quad f_u \cdot f_{vv} = F_v,$$
$$f_{uv} \cdot f_{uv} = \frac{G_{uu}}{2}, \quad f_{uv} \cdot f_{vv} = \frac{G_{uv}}{2}, \quad f_{vv} \cdot f_{vv} = \frac{G_{vv}}{2}$$

at $(u, v) = (0, 0)$. Since

$$|f_u|^2 = E, \quad |f_u \times f_{vv}|^2 = (f_u \cdot f_u)(f_{vv} \cdot f_{vv}) - (f_u \cdot f_{vv})^2 = \frac{EG_{vv}}{2} - (F_v)^2$$

at $(u, v) = (0, 0)$, we can conclude that a_{02} is an intrinsic invariant.

Similarly, to prove a_{20} and a_{11} are intrinsic, it is sufficient to show that $[f_u, f_{uu}, f_{vv}]$ and $[f_u, f_{uv}, f_{uu}]$ are both written in terms of derivatives of E, F and G at $(0, 0)$. In fact, (14) implies that $[f_u, f_{uu}, f_{vv}]$ is intrinsic, because

$$\begin{aligned}
[f_u, f_{uu}, f_{vv}] &= \frac{1}{[f_u, f_{uv}, f_{vv}]} \det \left(\begin{pmatrix} f_u \\ f_{uv} \\ f_{vv} \end{pmatrix} (f_u, f_{uu}, f_{vv}) \right) \\
&= \frac{1}{[f_u, f_{uv}, f_{vv}]} \det \begin{pmatrix} f_u \cdot f_u & f_u \cdot f_{uu} & f_u \cdot f_{vv} \\ f_{uv} \cdot f_u & f_{uv} \cdot f_{uu} & f_{uv} \cdot f_{vv} \\ f_{vv} \cdot f_u & f_{vv} \cdot f_{uu} & f_{vv} \cdot f_{vv} \end{pmatrix} \\
&= \frac{1}{[f_u, f_{uv}, f_{vv}]} \det \begin{pmatrix} E & f_u \cdot f_{uu} & F_v \\ F_u & f_{uv} \cdot f_{uu} & G_{uv}/2 \\ F_v & f_{vv} \cdot f_{uu} & G_{vv}/2 \end{pmatrix}
\end{aligned}$$

holds at $(u, v) = (0, 0)$, and

$$\begin{aligned}
f_u(0,0) \cdot f_{uu}(0,0) &= \frac{E_u(0,0)}{2}, \\
f_{uv}(0,0) \cdot f_{uu}(0,0) &= F_{uu}(0,0) - \frac{E_{uv}(0,0)}{2}, \\
f_{vv}(0,0) \cdot f_{uu}(0,0) &= F_{uv}(0,0) - \frac{E_{vv}(0,0)}{2}.
\end{aligned}$$

Similarly, $[f_u, f_{uv}, f_{uu}]$ is intrinsic, because of the identity

$$[f_u, f_{uv}, f_{uu}] = \frac{1}{[f_u, f_{uv}, f_{vv}]} \det \left(\begin{pmatrix} f_u \\ f_{uv} \\ f_{vv} \end{pmatrix} (f_u, f_{uv}, f_{uu}) \right).$$

□

Remark 7. The value $\Delta := [f_u(0,0), f_{uv}(0,0), f_{vv}(0,0)]$ is a criterion of cross cap singularities. In the above proof (cf. (14)), we showed the identity

$$\Delta^2 = \det \begin{pmatrix} E & F_u & F_v \\ F_u & G_{uu}/2 & G_{uv}/2 \\ F_v & G_{uv}/2 & G_{vv}/2 \end{pmatrix}$$

at $(u,v)=(0,0)$, which implies that Δ is intrinsic. Moreover, we set

$$h(u,v):=E(u,v)G(u,v)-(F(u,v))^2.$$

Using the fact that (u,v) is admissible, one can easily prove

$$\Delta^2=\frac{1}{4E(0,0)}\biggl(h_{uu}(0,0)h_{vv}(0,0)-(h_{uv}(0,0))^2\biggr),$$

that is, Δ is closely related to the Hessian of h. Since $h(u,v)$ is non-negative and $h(0,0)=0$, it holds that $h_{vv}(0,0)\geq 0$. Moreover, the identity

$$a_{02}=\frac{\sqrt{E(0,0)}(h_{vv}(0,0))^{3/2}}{2\Delta^2}$$

holds.

In [12], [11] and [3], ellipticity, hyperbolicity and parabolicity of cross caps are defined. The following assertion holds:

Corollary 8. *The ellipticity, hyperbolicity and parabolicity of cross caps in* $\boldsymbol{R}^3$ *are all intrinsic properties.*

Proof. A cross cap is elliptic (resp. hyperbolic) if $a_{20}>0$ (resp. $a_{20}<0$). Since we have already seen that a_{20} is intrinsic, ellipticity and hyperbolicity are as well. In [3], it was shown that a cross cap is parabolic if and only if $a_{20}=0$ and the zero set Z_K of the Gaussian curvature gives a regular curve in the $r\theta$-plane which is tangent to the line $r=0$, where

$$u=r\cos\theta,\qquad v=r\sin\theta$$

and (u,v) is a canonical coordinate system. Since the set Z_K is intrinsic and this tangency property does not depend on the choice of an admissible coordinate system, we get the assertion. $\Box$

We fix a cross cap $f:(U;u,v)\to\boldsymbol{R}^3$, where (u,v) is an admissible coordinate system, that is, $(u,v)=(0,0)$ is a cross cap singularity and $f_v(0,0)=0$. We call the line

$$\{f(0,0)+tf_u(0,0)\,;\,t\in\boldsymbol{R}\}$$

the *tangential line* at the cross cap and a non-zero vector at $T_{f(0,0)}\boldsymbol{R}^3$ proportional to $f_u(0,0)$ is called the *tangential direction*. The plane passing through $f(0,0)$ spanned by $f_u(0,0)$ and $f_{vv}(0,0)$ is called the *principal plane*. On the other hand, the plane passing through $f(0,0)$ perpendicular to the proper tangential direction is called the *normal plane*. The unit normal vector $\nu(u,v)$ near the cross cap at $(u,v)=(0,0)$ can be extended as a C^∞-function of r, θ by setting $u=r\cos\theta$ and $v=r\sin\theta$, and the limiting normal vector

$$\nu(\theta):=\lim_{r\to 0}\nu(r\cos\theta,r\sin\theta)\in T_{f(0,0)}\boldsymbol{R}^3$$

lies in the normal plane. Then, one can consider the parallel family of cross caps

$$f_t(r,\theta):=f(r,\theta)+t\nu(r,\theta) \tag{15}$$

which are C^∞-maps with respect to (r,θ), even at $r=0$. The focal surface of this parallel family meets the normal plane at the focal conic as mentioned in the introduction. On the other hand, the principal plane has the following property:

Proposition 9. *The initial velocity vector of the space curve emanating from the cross cap singularity which parametrizes the self-intersection is contained in the principal plane.*

Proof. Since the principal plane is invariant under diffeomorphisms of $\boldsymbol{R}^3$, this assertion can be verified by the standard cross cap. $\square$

As pointed out in the introduction, a_{20} is an important intrinsic invariant of cross caps related to the sign of the Gaussian curvature. The following assertion can be proved easily, which gives a geometric meaning for a_{20}:

Proposition 10. *The section of a cross cap by its principal plane contains a regular curve γ whose velocity vector is $f_u(0,0)$, and then the curvature of γ as a plane curve at the cross cap is equal to a_{20}, where we give the orientation to the principal plane so that $\{f_u(0,0), f_{vv}(0,0)\}$ is a positive frame.*

Intersections of a cross cap with planes are discussed in [4].

Finally, we discuss on cross caps in an arbitrary Riemannian 3-manifold (N^3,g). Let $f: M^2 \to (N^3,g)$ be a C^∞-map having a cross cap singularity at $p \in M^2$, where M^2 is a 2-manifold. Then there exists a local coordinate system (u,v) of M^2 centered at p and a normal coordinate system (x,y,z) of (N^3,g) centered at $f(p)$ such that (cf. (2))

$$f(u,v) = \Big(u, uv, \frac{a_{20}}{2}u^2 + a_{11}uv + \frac{a_{02}}{2}v^2\Big) + O(u,v)^3.$$

Like as in the case of the Euclidean 3-space, one can easily verify that a_{20}, a_{02} and a_{11} are all intrinsic invariants: In fact, we set

$$[\boldsymbol{a},\boldsymbol{b},\boldsymbol{c}] := \Omega(\boldsymbol{a},\boldsymbol{b},\boldsymbol{c}) = g(\boldsymbol{a}\times\boldsymbol{b},\boldsymbol{c}),$$

where Ω is the Riemannian volume form of (N^3,g) and $\boldsymbol{a},\boldsymbol{b},\boldsymbol{c}$ are vector fields of N^3 along the C^∞-map f. We denote by D the Levi-Civita connection of g. By replacing

$$f_{uu} \mapsto D_u f_u, \quad f_{uv} \mapsto D_u f_v, \quad f_{vv} \mapsto D_v f_v,$$

the three formulas (11), (12) and (13) hold at the cross cap singularity of N^3.

By a straightforward calculation, the first and the second fundamental forms

$$E\,du^2 + 2F\,du\,dv + G\,dv^2, \quad L\,du^2 + 2M\,du\,dv + N\,dv^2$$

have the following expressions

$$
\begin{aligned}
&E = 1 + r^2\big(\sin^2\theta + (a_{20}\cos\theta + a_{11}\sin\theta)^2 + O(r)\big),\\
&F = r^2\left(a_{20}a_{11}\cos^2\theta + (1 + a_{20}a_{02} + a_{11}^2)\sin\theta\cos\theta + a_{11}a_{02}\sin^2\theta + O(r)\right),\\
&G = r^2\big(A_\theta^2 + O(r)\big),\\
&L = \frac{a_{20}\cos\theta}{A_\theta} + O(r), \quad M = -\frac{a_{02}\sin\theta}{A_\theta} + O(r), \quad N = \frac{a_{02}\cos\theta}{A_\theta} + O(r),
\end{aligned}
$$

where $u = r\cos\theta$, $v = r\sin\theta$ and

$$A_\theta := \sqrt{\cos^2\theta + (a_{11}\cos\theta + a_{02}\sin\theta)^2}.$$

We denote by K_{ext} the determinant of the shape operator of f, which is called the *extrinsic curvature function*.

Since $EG-F^2 = r^2(A_\theta^2+O(r))$, the mean curvature function H, the extrinsic curvature function K_{ext}, and the Gaussian curvature function K are given by

$$H = \frac{1}{r^2}\left(\frac{a_{02}\cos\theta}{2A_\theta^3} + O(r)\right), \tag{16}$$

$$K_{\text{ext}} = \frac{a_{02}}{r^2A_\theta^4}\left(a_{20}\cos^2\theta - a_{02}\sin^2\theta + O(r)\right), \tag{17}$$

$$K = K_{\text{ext}} + c_g(r,\theta) = \frac{a_{02}}{r^2A_\theta^4}\left(a_{20}\cos^2\theta - a_{02}\sin^2\theta + O(r)\right), \tag{18}$$

where $c_g(r,\theta)$ is a suitable C^∞-function at p with respect to the sectional curvature of the Riemannian metric g appeared in the Gauss equation. When (N^3, g) is the Euclidean space, the formulas (16) and (17) (and also the description of principal curvatures) have been given in [3]. It should be remarked that the top terms of the curvature functions K and H are determined by the three invariants a_{20}, a_{02} and a_{11}. This fact seems a remarkable property of cross caps since *the top terms of H and K do not depend on a choice of ambient spaces*. Moreover, (17) and (18) imply that the asymptotic behaviors of K and K_{ext} are same at the cross cap singularity.

The following is a generalization of the assertion proved in Fukui-Ballesteros [2] and Tari [11] when (N^3, g) is the Euclidean 3-space:

Proposition 11. *Umbilical points do not accumulate to a cross cap in (N^3, g).*

Proof. By (17), we know that $K_{\text{ext}} < 0$ if $\theta = \pm\pi/2$. Thus it is sufficient to show that $H^2 - K_{\text{ext}}$ does not vanish under the assumption $\cos\theta \neq 0$. In fact

$$H^2 - K_{\text{ext}} = \frac{1}{r^4}\left(\frac{(a_{02}\cos\theta)^2}{4A_\theta^6} + O(r)\right)$$

diverges if $\cos\theta \neq 0$ as r tends to zero. □

As noted in the introduction, the ellipticity and hyperbolicity of cross caps are determined by the sign of the invariant a_{20}. In this paper, we have shown the existence of non-trivial isometric deformations of quadratic cross caps when a_{20}

vanishes. We end this paper with the following rigidity problem for generic cross caps in $\boldsymbol{R}^3$:

Question. *Supposing that two cross caps in* $\boldsymbol{R}^3$ *have a common first fundamental form and satisfy a suitable genericity assumption, would they then be congruent?*

Acknowledgments

The authors thank Shyuichi Izumiya, Toshizumi Fukui and Wayne Rossman for valuable comments.

References

[1] J. W. Bruce and J. M. West, *Functions on cross-caps*, Math. Proc. Cambridge Philos. Soc., **123** (1998), 19–39.
[2] T. Fukui, J. J. Nuño-Ballesteros, *Isolated roundings and flattenings of submanifolds in Euclidean spaces*, Tôhoku Math. J., **57** (2005), 469–503.
[3] T. Fukui and M. Hasegawa, *Fronts of Whitney umbrella— a differential geometric approach via blowing up—*, J. Singularities, **4** (2012), 35–67.
[4] T. Fukui and M. Hasegawa, *Height functions on Whitney umbrellas*, preprint, arXiv:1205.3044.
[5] R. Garcia, C. Gutierrez and J. Sotomayor, *Lines of principal curvature around umbilics and Whitney umbrellas*, Tôhoku Math. J., **52** (2000), 163–172.
[6] C. Gutierrez and J. Sotomayor, *Lines of principal curvature for mappings with Whitney umbrella singularities*, Tôhoku Math. J., **38** (1986), 551–559.
[7] N. H. Kuiper, *Stable surfaces in Euclidean three space*, Math. Scand., **36** (1975), 83–96.
[8] J. J. Nuño-Ballesteros and F. Tari, *Surfaces in* $\boldsymbol{R}^4$ *and their projections to* 3*-spaces*, Proc. Roy. Soc. Edinburgh Sect. A, **137** (2007), 1313–1328.
[9] K. Saji, M. Umehara and K. Yamada, *The geometry of fronts*, Ann. of Math., **169** (2009) 491–529.
[10] J. M. Oliver, *On pairs of foliations of a parabolic cross-cap*, Qual. Theory Dyn. Syst., **10** (2011), 139–166.
[11] F. Tari, *Pairs of geometric foliations on a cross-cap*, Tôhoku Math. J., **59** (2007), 233–258.
[12] J. M. West, *The differential geometry of the cross-cap*, Ph. D. thesis, The University of Liverpool, 1995.

(Masaru Hasegawa) Department of Mathematics, Faculty of Science, Saitama University, Sakura-ku, Saitama 338-857, Japan
E-mail address: `mhasegawa@mail.saitama-u.ac.jp`

(MasaakiUmehara) Department of Mathematical and Computing Sciences, Tokyo Institute of Technology, 2-12-1-W8-34, O-okayama Meguro-ku, Tokyo 152-8552 Japan
E-mail address: `umehara@is.titech.ac.jp`

(Atsufumi Honda, Kosuke Naokawa and Kotaro Yamada) Department of Mathematics, Tokyo Institute of Technology, O-okayama, Meguro, Tokyo 152-8551, Japan
E-mail address: `10d00059@math.titech.ac.jp`
E-mail address: `naokawa1@is.titech.ac.jp`
E-mail address: `kotaro@math.titech.ac.jp`